

## QuantiGene® Bibliography

### 2009

1. Berrodin, T.J., et al., Novel progesterone receptor modulators with gene selective and context-dependent partial agonism. *Biochem Pharmacol*, 2009. 77(2): p. 204-15.
2. Buckley, D.B. and C.D. Klaassen, Induction of mouse UDP-glucuronosyltransferase mRNA expression in liver and intestine by activators of aryl-hydrocarbon receptor, constitutive androstane receptor, pregnane X receptor, peroxisome proliferator-activated receptor alpha, and nuclear factor erythroid 2-related factor 2. *Drug Metab Dispos*, 2009. 37(4): p. 847-56.
3. Buckley, D.B. and C.D. Klaassen, Mechanism of gender-divergent UDP-glucuronosyltransferase mRNA expression in mouse liver and kidney. *Drug Metab Dispos*, 2009. 37(4): p. 834-40.
4. Champion, S.N., et al., Effect of allyl alcohol on hepatic transporter expression: Zonal patterns of expression and role of Kupffer cell function. *Toxicology and Applied Pharmacology*, 2009. 236: p. 49–58.
5. Cui, Y.J., et al., Tissue distribution, gender-divergent expression, ontogeny, and chemical induction of multidrug resistance transporter genes (Mdr1a, Mdr1b, Mdr2) in mice. *Drug Metab Dispos*, 2009. 37(1): p. 203-10.
6. Judge, A.D., et al., Confirming the RNAi-mediated mechanism of action of siRNA-based cancer therapeutics in mice. *J Clin Invest*, 2009. 119(3): p. 661-73.
7. Mahmoudi, S., et al., Wrap53, a natural p53 antisense transcript required for p53 induction upon DNA damage. *Mol Cell*, 2009. 33(4): p. 462-71.
8. Reisman, S.A., L.M. Aleksunes, and C.D. Klaassen, Oleanolic acid activates Nrf2 and protects from acetaminophen hepatotoxicity via Nrf2-dependent and Nrf2-independent processes. *Biochem Pharmacol*, 2009. 77(7): p. 1273-82.
9. Reisman, S.A., et al., CDDO-Im protects from acetaminophen hepatotoxicity through induction of Nrf2-dependent genes. *Toxicology and Applied Pharmacology*, 2009. 236: p. 109–114.
10. Shinonaga, H., et al., Pochonins K–P: new radicicol analogues from *Pochonia chlamydosporia* var. *chlamydosporia* and their WNT-5A expression inhibitory activities. *Tetrahedron*, 2009. 65: p. 3446–3453.
11. Shinonaga, H., A.I. Yoji Kawamura, Mari Aoki, Noriyoshi Sakai, and A.K. Natsuko Fujimoto, The search for a hair-growth stimulant: new radicicol analogues as WNT-5A expression inhibitors from *Pochonia chlamydosporia* var. *chlamydosporia*. *Tetrahedron Letters*, 2009. 50: p. 108-110.
12. Vickers, A.E., Characterization of hepatic mitochondrial injury induced by fatty acid oxidation inhibitors. *Toxicol Pathol*, 2009. 37(1): p. 78-88.
13. Zhang, Y.K., R.L. Yeager, and C.D. Klaassen, Circadian expression profiles of drug-processing genes and transcription factors in mouse liver. *Drug Metab Dispos*, 2009. 37(1): p. 106-15.
14. Zhu, J., et al., Regulation of apolipoprotein AI (apoAI) processing by Procollagen C-proteinase enhancer (PCPE2) and bone morphogenetic protein-1 (BMP-1). *J Lipid Res*, 2009.

### 2008

1. Aagaard, L.A., et al., Engineering and optimization of the miR-106b cluster for ectopic expression of multiplexed anti-HIV RNAs. *Gene Ther*, 2008. 15(23): p. 1536-49.
2. Akinc, A., et al., A combinatorial library of lipid-like materials for delivery of RNAi therapeutics. *Nat Biotechnol*, 2008. 26(5): p. 561-9.
3. Aleksunes, L.M., et al., Induction of Mrp3 and Mrp4 transporters during acetaminophen hepatotoxicity is dependent on Nrf2. *Toxicol Appl Pharmacol*, 2008. 226(1): p. 74-83.
4. Aleksunes, L.M., et al., Renal xenobiotic transporters are differentially expressed in mice following cisplatin treatment. *Toxicology*, 2008. 250(2-3): p. 82-8.
5. Aleksunes, L.M., Y. Cui, and C.D. Klaassen, Prominent Expression of Xenobiotic Efflux Transporters in Mouse Extraembryonic Fetal Membranes Compared to Placenta. *Drug Metab Dispos*, 2008.

6. Aleku, M., et al., Atu027, a liposomal small interfering RNA formulation targeting protein kinase N3, inhibits cancer progression. *Cancer Res*, 2008. 68(23): p. 9788-98.
7. Alnouti, Y. and C.D. Klaassen, Regulation of sulfotransferase enzymes by prototypical microsomal enzyme inducers in mice. *J Pharmacol Exp Ther*, 2008. 324(2): p. 612-21.
8. Alnouti, Y. and C.D. Klaassen, Tissue distribution, ontogeny, and regulation of aldehyde dehydrogenase (aldh) enzymes mRNA by prototypical microsomal enzyme inducers in mice. *Toxicol Sci*, 2008. 101(1): p. 51-64.
9. Anderson, E.M., et al., Experimental validation of the importance of seed complement frequency to siRNA specificity. *Rna*, 2008. 14(5): p. 853-61.
10. Arikawa, E., et al., Cross-platform comparison of SYBR Green real-time PCR with TaqMan PCR, microarrays and other gene expression measurement technologies evaluated in the MicroArray Quality Control (MAQC) study. *BMC Genomics*, 2008. 9: p. 328.
11. Beilke, L.D., et al., Minimal Role of Hepatic Transporters in the Hepatoprotection against LCA-Induced Intrahepatic Cholestasis. *Toxicol Sci*, 2008. 102(1): p. 196-204.
12. Calcagno, A.M., et al., Single-step doxorubicin-selected cancer cells overexpress the ABCG2 drug transporter through epigenetic changes. *Br J Cancer*, 2008. 98(9): p. 1515-24.
13. Chae, Y.K., et al., Human AQP5 plays a role in the progression of chronic myelogenous leukemia (CML). *PLoS ONE*, 2008. 3(7): p. e2594.
14. Cheng, X. and C.D. Klaassen, Perfluorocarboxylic acids induce cytochrome P450 enzymes in mouse liver through activation of PPAR-alpha and CAR transcription factors. *Toxicol Sci*, 2008. 106(1): p. 29-36.
15. Cheng, X. and C.D. Klaassen, Critical role of PPAR-alpha in perfluorooctanoic acid- and perfluorodecanoic acid-induced downregulation of Oatp uptake transporters in mouse livers. *Toxicol Sci*, 2008. 106(1): p. 37-45.
16. Chung, J.Y., et al., Factors in tissue handling and processing that impact RNA obtained from formalin-fixed, paraffin-embedded tissue. *J Histochem Cytochem*, 2008. 56(11): p. 1033-42.
17. Crombez, L., et al., A New Potent Secondary Amphipathic Cell-penetrating Peptide for siRNA Delivery Into Mammalian Cells. *Mol Ther*, 2008.
18. Eastham, L.L., C.N. Mills, and R.M. Niles, PPARalpha/gamma Expression and Activity in Mouse and Human Melanocytes and Melanoma Cells. *Pharm Res*, 2008.
19. Frank-Kamenetsky, M., et al., Therapeutic RNAi targeting PCSK9 acutely lowers plasma cholesterol in rodents and LDL cholesterol in nonhuman primates. *Proc Natl Acad Sci U S A*, 2008.
20. Grueneberg, D.A., et al., Kinase requirements in human cells: IV. Differential kinase requirements in cervical and renal human tumor cell lines. *Proc Natl Acad Sci U S A*, 2008. 105(43): p. 16490-5.
21. Gyamfi, M.A., et al., Hepatic effects of a methionine-choline-deficient diet in hepatocyte RXRalpha-null mice. *Toxicol Appl Pharmacol*, 2008.
22. Haag, J., et al., Efficient non-viral transfection of primary human adult chondrocytes in a high-throughput format. *Osteoarthritis Cartilage*, 2008.
23. Hart, S.N., et al., Genetic polymorphisms in cytochrome P450 oxidoreductase influence microsomal P450-catalyzed drug metabolism. *Pharmacogenet Genomics*, 2008. 18(1): p. 11-24.
24. Halpern, M.D. and B. Dvorak, Does Abnormal Bile Acid Metabolism Contribute to NEC? *Semin Perinatol*, 2008. 32(2): p. 114-21.
25. Honda, M., et al., Specific knockdown of m-calpain blocks myogenesis with cDNA deduced from the corresponding RNAi. *Am J Physiol Cell Physiol*, 2008.
26. Hubbard, K., et al., Knockdown of TFIIS by RNA silencing inhibits cancer cell proliferation and induces apoptosis. *BMC Cancer*, 2008. 8: p. 133.
27. Jonson, A.L., et al., Gene silencing with siRNA targeting E6/E7 as a therapeutic intervention in a mouse model of cervical cancer. *Gynecol Oncol*, 2008.

28. Knudsen, B.S., et al., Evaluation of the Branched-Chain DNA Assay for Measurement of RNA in Formalin-Fixed Tissues. *J Mol Diagn*, 2008.
29. Lee, A.C., et al., Electrochemical Branched-DNA Assay for Polymerase Chain Reaction-Free Detection and Quantification of Oncogenes in Messenger RNA. *Anal Chem*, 2008.
30. Lorenzi, P.L., et al., Asparagine synthetase is a predictive biomarker of L-asparaginase activity in ovarian cancer cell lines. *Mol Cancer Ther*, 2008. 7(10): p. 3123-8.
31. Lu, H. and C. Klaassen, Gender differences in mRNA expression of ATP-binding cassette efflux and bile acid transporters in kidney, liver, and intestine of 5/6 nephrectomized rats. *Drug Metab Dispos*, 2008. 36(1): p. 16-23.
32. Lu, H., et al., Characterization of Organic Anion Transporting Polypeptide 1b2-null Mice: Essential Role in Hepatic Uptake/Toxicity of Phalloidin and Microcystin-LR. *Toxicol Sci*, 2008.
33. Maher, J.M., et al., Nrf2- and PPAR alpha-mediated regulation of hepatic Mrp transporters after exposure to perfluorooctanoic acid and perfluorodecanoic acid. *Toxicol Sci*, 2008. 106(2): p. 319-28.
34. Merrell, M.D., et al., The Nrf2 activator oltipraz also activates the constitutive androstane receptor. *Drug Metab Dispos*, 2008. 36(8): p. 1716-21.
35. Migliaccio, C.T., et al., The IL-4Ralpha pathway in macrophages and its potential role in silica-induced pulmonary fibrosis. *J Leukoc Biol*, 2008. 83(3): p. 630-9.
36. Moore, M., et al., Measurement of neutralising antibodies to type I interferons by gene expression assays specific for type 1 interferon-inducible 6-16 mRNA. *J Pharm Biomed Anal*, 2008.
37. Nair, A., et al., A rapid and efficient branched DNA hybridization assay to titer lentiviral vectors. *J Virol Methods*, 2008. 153(2): p. 269-72.
38. Querbes, W., et al., Direct CNS Delivery of siRNA Mediates Robust Silencing in Oligodendrocytes. *Oligonucleotides*, 2008.
39. Robbins, M., et al., Misinterpreting the therapeutic effects of siRNA caused by immune stimulation. *Hum Gene Ther*, 2008.
40. Ryan, M.C., et al., SpliceCenter: a suite of web-based bioinformatic applications for evaluating the impact of alternative splicing on RT-PCR, RNAi, microarray, and peptide-based studies. *BMC Bioinformatics*, 2008. 9: p. 313.
41. Sano, M., et al., Expression of Long Anti-HIV-1 Hairpin RNAs for the Generation of Multiple siRNAs: Advantages and Limitations. *Mol Ther*, 2008. 16(1): p. 170-7.
42. Shou, M., et al., Modeling, prediction, and in vitro in vivo correlation of CYP3A4 induction. *Drug Metab Dispos*, 2008. 36(11): p. 2355-70.
43. Simpson, K.J., et al., Identification of genes that regulate epithelial cell migration using an siRNA screening approach. *Nat Cell Biol*, 2008.
44. Tanaka, Y., et al., Nf-E2-Related Factor 2 (Nrf2) Inhibits Lipid Accumulation and Oxidative Stress in Mice Fed a High-Fat Diet. *J Pharmacol Exp Ther*, 2008.
45. Tanaka, Y., et al., Coordinated induction of Nrf2 target genes protects against iron nitrilotriacetate (FeNTA)-induced nephrotoxicity. *Toxicol Appl Pharmacol*, 2008. 231(3): p. 364-73.
46. Tanaka, Y., et al., Ischemia-reperfusion of rat livers decreases liver and increases kidney multidrug resistance associated protein 2 (Mrp2). *Toxicol Sci*, 2008. 101(1): p. 171-8.
47. Unwalla, H.J., et al., Use of a U16 snoRNA-containing ribozyme library to identify ribozyme targets in HIV-1. *Mol Ther*, 2008. 16(6): p. 1113-9.
48. Zhang, Y.K., R.L. Yeager, and C.D. Klaassen, Circadian Expression Profiles of Drug Processing Genes and Transcription Factors in Mouse Liver. *Drug Metab Dispos*, 2008.

## 2007

1. Aleksunes, L.M., et al., Influence of acetaminophen vehicle on regulation of transporter gene expression during hepatotoxicity. *J Toxicol Environ Health A*, 2007. 70(21): p. 1870-2.

2. Barnes, S.N., et al., Induction of hepatobiliary efflux transporters in acetaminophen-induced acute liver failure cases. *Drug Metab Dispos*, 2007. 35(10): p. 1963-9.
3. Buckley, D.B. and C.D. Klaassen, Tissue- and gender-specific mRNA expression of UDP-glucuronosyltransferases (UGTs) in mice. *Drug Metab Dispos*, 2007. 35(1): p. 121-7.
4. Chen, C., et al., Activation of cAMP-dependent signaling pathway induces mouse organic anion transporting polypeptide 2 expression. *Mol Pharmacol*, 2007. 71(4): p. 1159-64.
5. Cheshenko, K., et al., Expression of Zebra Fish Aromatase cyp19a and cyp19b Genes in Response to the Ligands of Estrogen Receptor and Aryl Hydrocarbon Receptor. *Toxicol Sci*, 2007. 96(2): p. 255-67.
6. Coates, S.S., et al., Beryllium induces premature senescence in human fibroblasts. *J Pharmacol Exp Ther*, 2007. 322(1): p. 70-9.
7. Dotzlaw, J., et al., Derivation and characterization of monoclonal antibodies against human folypolyglutamate synthetase. *Hybridoma (Larchmt)*, 2007. 26(3): p. 155-61.
8. Fisher, C.D., et al., Induction of drug-metabolizing enzymes by garlic and allyl sulfide compounds via activation of constitutive androstane receptor and nuclear factor E2-related factor 2. *Drug Metab Dispos*, 2007. 35(6): p. 995-1000.
9. Hernandez, S., B. Chavez Munguia, and L. Gonzalez-Mariscal, ZO-2 silencing in epithelial cells perturbs the gate and fence function of tight junctions and leads to an atypical monolayer architecture. *Exp Cell Res*, 2007. 313(8): p. 1533-47.
10. John, M., et al., Effective RNAi-mediated gene silencing without interruption of the endogenous microRNA pathway. *Nature*, 2007. 449(7163): p. 745-7.
11. Knight, T.R., S. Choudhuri, and C.D. Klaassen, Constitutive mRNA expression of various glutathione S-transferase isoforms in different tissues of mice. *Toxicol Sci*, 2007. 100(2): p. 513-24.
12. Lantz, R.C., et al., The effect of extracts from ginger rhizome on inflammatory mediator production. *Phytomedicine*, 2007. 14(2-3): p. 123-8.
13. Lickteig, A.J., et al., Genes of the antioxidant response undergo upregulation in a rodent model of nonalcoholic steatohepatitis. *J Biochem Mol Toxicol*, 2007. 21(4): p. 216-20.
14. Lickteig, A.J., et al., Efflux transporter expression and acetaminophen metabolite excretion are altered in rodent models of nonalcoholic fatty liver disease. *Drug Metab Dispos*, 2007. 35(10): p. 1970-8.
15. Lickteig, A.J., et al., Differential regulation of hepatic transporters in the absence of tumor necrosis factor-alpha, interleukin-1beta, interleukin-6, and nuclear factor-kappaB in two models of cholestasis. *Drug Metab Dispos*, 2007. 35(3): p. 402-9.
16. Lin, X., et al., 'Seed' analysis of off-target siRNAs reveals an essential role of Mcl-1 in resistance to the small-molecule Bcl-2/Bcl-XL inhibitor ABT-737. *Oncogene*, 2007. 26(27): p. 3972-9.
17. Martin, S.E., et al., Multiplexing siRNAs to compress RNAi-based screen size in human cells. *Nucleic Acids Res*, 2007. doi:10.1093/nar/gkm141.
18. Miao, Z., et al., CXCR7 (RDC1) promotes breast and lung tumor growth in vivo and is expressed on tumor-associated vasculature. *Proc Natl Acad Sci U S A*, 2007. 104(40): p. 15735-40.
19. Miao, Z.H., et al., Nonclassic functions of human topoisomerase I: genome-wide and pharmacologic analyses. *Cancer Res*, 2007. 67(18): p. 8752-61.
20. Pacyniak, E.K., et al., The Flame Retardants, Polybrominated Diphenyl Ethers (PBDE), are Pregnane X Receptor (PXR) Activators. *Toxicol Sci*, 2007.
21. Petrick, J.S. and C.D. Klaassen, Importance of hepatic induction of constitutive androstane receptor and other transcription factors that regulate xenobiotic metabolism and transport. *Drug Metab Dispos*, 2007. 35(10): p. 1806-15.
22. Slitt, A.L., et al., Regulation of transporter expression in mouse liver, kidney, and intestine during extrahepatic cholestasis. *Biochim Biophys Acta*, 2007. 1768(3): p. 637-47.
23. Tanaka, Y., et al., Hepatic ischemia-reperfusion induces renal heme oxygenase-1 via NF-E2-related factor 2 in rats and mice. *Mol Pharmacol*, 2007. 71(3): p. 817-25.

24. Vermeulen, A., et al., Double-stranded regions are essential design components of potent inhibitors of RISC function. *RNA*, 2007. 13(5): p. 723-30.
25. Wei, X., et al., Development and validation of a quantitative cell-based bioassay for comparing the pharmacokinetic profiles of two recombinant erythropoietic proteins in serum. *J Pharm Biomed Anal*, 2007. 43(2): p. 666-76.
26. Wen, X. and T. Walle, Cytochrome P450 1B1, a novel chemopreventive target for benzo[a]pyrene-initiated human esophageal cancer. *Cancer Lett*, 2007. 246(1-2): p. 109-14.
27. Wolfrum, C., et al., Mechanisms and optimization of in vivo delivery of lipophilic siRNAs. *Nat Biotechnol*, 2007. 25(10): p. 1149-57.
28. Yamaguchi, T., et al., Identification of JTP-70902, a p15(INK4b)-inductive compound, as a novel MEK1/2 inhibitor. *Cancer Sci*, 2007. 98(11): p. 1809-16.
29. Yuan, Q., et al., Age-dependent expression of the cystatin-related epididymal spermatogenic (Cres) gene in mouse testis and epididymis. *Asian J Androl*, 2007. 9(3): p. 305-11.
30. Zhang, X., N.J. Cherrington, and S.H. Wright, Molecular identification and functional characterization of rabbit MATE1 and MATE2-K. *Am J Physiol Renal Physiol*, 2007. 293(1): p. F360-70.
31. Zhou, Y., et al., Phenotypic and structural analyses of hepatitis C virus NS3 protease Arg155 variants: sensitivity to telaprevir (VX-950) and interferon alpha. *J Biol Chem*, 2007. 282(31): p. 22619-28.

## 2006

1. Aleksunes, L.M., et al., Nuclear factor-E2-related factor 2 expression in liver is critical for induction of NAD(P)H:quinone oxidoreductase 1 during cholestasis. *Cell Stress Chaperones*, 2006. 11(4): p. 356-63.
2. Alnouti, Y. and C.D. Klaassen, Tissue distribution and ontogeny of sulfotransferase enzymes in mice. *Toxicol Sci*, 2006. 93(2): p. 242-55.
3. Alnouti, Y., J.S. Petrick, and C.D. Klaassen, Tissue distribution and ontogeny of organic cation transporters in mice. *Drug Metab Dispos*, 2006. 34(3): p. 477-82.
4. Canales, R.D., et al., Evaluation of DNA microarray results with quantitative gene expression platforms. *Nat Biotechnol*, 2006. 24(9): p. 1115-22.
5. Cheng, X. and C.D. Klaassen, Regulation of mRNA expression of xenobiotic transporters by the pregnane x receptor in mouse liver, kidney, and intestine. *Drug Metab Dispos*, 2006. 34(11): p. 1863-7.
6. Cheng, X., et al., Endocrine regulation of gender-divergent mouse organic anion-transporting polypeptide (Oatp) expression. *Mol Pharmacol*, 2006. 70(4): p. 1291-7.
7. Conrad, P.G., R.V. Chavli, and R.S. Givens, Caged substrates applied to high content screening: an introduction with an eye to the future. *Methods Mol Biol*, 2006. 356: p. 253-66.
8. Das, P.C., et al., Fipronil induces CYP isoforms and cytotoxicity in human hepatocytes. *Chem Biol Interact*, 2006. 164(3): p. 200-14.
9. Deng, D.X., et al., Differences in vascular bed disease susceptibility reflect differences in gene expression response to atherogenic stimuli. *Circ Res*, 2006. 98(2): p. 200-8.
10. Dotzlaw, J., et al., Derivation and characterization of a monoclonal antibody against human glycinamide ribonucleotide formyltransferase. *Hybridoma (Larchmt)*, 2006. 25(3): p. 139-44.
11. Fedorov, Y., et al., Off-target effects by siRNA can induce toxic phenotype. *RNA*, 2006. 12(7): p. 1188-96.
12. Ferrari, N., et al., Characterization of antisense oligonucleotides comprising 2'-deoxy-2'-fluoro-beta-D-arabinonucleic acid (FANA): specificity, potency, and duration of activity. *Ann N Y Acad Sci*, 2006. 1082: p. 91-102.
13. Fortin, M., et al., Effects of antisense oligodeoxynucleotides targeting CCR3 on the airway response to antigen in rats. *Oligonucleotides*, 2006. 16(3): p. 203-12.
14. Groves, C.E., et al., Sex differences in the mRNA, protein, and functional expression of organic anion transporter (Oat) 1, Oat3, and organic cation transporter (Oct) 2 in rabbit renal proximal tubules. *J Pharmacol Exp Ther*, 2006. 316(2): p. 743-52.

15. Hinfray, N., et al., Brain and gonadal aromatase as potential targets of endocrine disrupting chemicals in a model species, the zebrafish (*Danio rerio*). *Environ Toxicol*, 2006. 21(4): p. 332-7.
16. Judge, A.D., et al., Design of noninflammatory synthetic siRNA mediating potent gene silencing in vivo. *Mol Ther*, 2006. 13(3): p. 494-505.
17. Kolachala, V.L., et al., Adenosine 2b receptor (A2bR) signals through adenylate cyclase (AC) 6 isoform in the intestinal epithelial cells. *Biochim Biophys Acta*, 2006. 1760(7): p. 1102-8.
18. Laville, N., et al., Modulation of aromatase activity and mRNA by various selected pesticides in the human choriocarcinoma JEG-3 cell line. *Toxicology*, 2006. 228(1): p. 98-108.
19. Lin, X., et al., 'Seed' analysis of off-target siRNAs reveals an essential role of Mcl-1 in resistance to the small-molecule Bcl-2/Bcl-X(L) inhibitor ABT-737. *Oncogene*, 2006.
20. Lorenzi, P.L., et al., Asparagine synthetase as a causal, predictive biomarker for L-asparaginase activity in ovarian cancer cells. *Mol Cancer Ther*, 2006. 5(11): p. 2613-2623.
21. Lu, H., X. Lei, and C. Klaassen, Gender differences in renal nuclear receptors and aryl hydrocarbon receptor in 5/6 nephrectomized rats. *Kidney Int*, 2006. 70(11): p. 1920-8.
22. Maher, J.M., et al., Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1alpha. *Biochem Pharmacol*, 2006. 72(4): p. 512-22.
23. Maher, J.M., et al., Hormonal regulation of renal multidrug resistance-associated proteins 3 and 4 (Mrp3 and Mrp4) in mice. *Biochem Pharmacol*, 2006. 71(10): p. 1470-8.
24. Majercak, J., et al., LRRTM3 promotes processing of amyloid-precursor protein by BACE1 and is a positional candidate gene for late-onset Alzheimer's disease. *Proc Natl Acad Sci U S A*, 2006. 103(47): p. 17967-17972.
25. Moffit, J.S., et al., Induction of hepatic transporters multidrug resistance-associated proteins (Mrp) 3 and 4 by clofibrate is regulated by peroxisome proliferator-activated receptor alpha. *J Pharmacol Exp Ther*, 2006. 317(2): p. 537-45.
26. Nguyen, Q.N., et al., Light controllable siRNAs regulate gene suppression and phenotypes in cells. *Biochim Biophys Acta*, 2006. 1758(3): p. 394-403.
27. Nishiu, J., et al., JTP-426467 acts as a selective antagonist for peroxisome proliferator-activated receptor gamma in vitro and in vivo. *Diabetes Obes Metab*, 2006. 8(5): p. 508-16.
28. Pruijboom-Brees, I., et al., A critical role for peroxisomal proliferator-activated receptor-alpha nuclear receptors in the development of cardiomyocyte degeneration and necrosis. *Am J Pathol*, 2006. 169(3): p. 750-60.
29. Reynolds, A., et al., Induction of the interferon response by siRNA is cell type- and duplex length-dependent. *RNA*, 2006. 12(6): p. 988-93.
30. Saetrom, P., et al., Conserved microRNA characteristics in mammals. *Oligonucleotides*, 2006. 16(2): p. 115-44.
31. Schmidt, R.J., et al., A 15-ketosterol is a liver X receptor ligand that suppresses sterol-responsive element binding protein-2 activity. *J Lipid Res*, 2006. 47(5): p. 1037-44.
32. Shelby, M.K. and C.D. Klaassen, Induction of rat UDP-glucuronosyltransferases in liver and duodenum by microsomal enzyme inducers that activate various transcriptional pathways. *Drug Metab Dispos*, 2006. 34(10): p. 1772-8.
33. Shi, L., et al., The MicroArray Quality Control (MAQC) project shows inter- and intraplatform reproducibility of gene expression measurements. *Nat Biotechnol*, 2006. 24(9): p. 1151-1161.
34. Slitt, A.L., et al., trans-Stilbene oxide induces expression of genes involved in metabolism and transport in mouse liver via CAR and Nrf2 transcription factors. *Mol Pharmacol*, 2006. 69(5): p. 1554-63.
35. Slitt, A.L., et al., Induction of genes for metabolism and transport by trans-stilbene oxide in livers of Sprague-Dawley and Wistar-Kyoto rats. *Drug Metab Dispos*, 2006. 34(7): p. 1190-7.
36. Stanley, L.A., et al., PXR and CAR: nuclear receptors which play a pivotal role in drug disposition and chemical toxicity. *Drug Metab Rev*, 2006. 38(3): p. 515-97.

37. Tanaka, Y., et al., Kupffer cell-mediated downregulation of hepatic transporter expression in rat hepatic ischemia-reperfusion. *Transplantation*, 2006. 82(2): p. 258-66.
38. Tsuji, P.A. and T. Walle, Inhibition of benzo[a]pyrene-activating enzymes and DNA binding in human bronchial epithelial BEAS-2B cells by methoxylated flavonoids. *Carcinogenesis*, 2006. 27(8): p. 1579-85.
39. Yang, W., et al., Direct quantification of gene expression in homogenates of formalin-fixed, paraffin-embedded tissues. *Biotechniques*, 2006. 40(4): p. 481-6.
40. Yu, M., J. Yeh, and C. Van Waes, Protein kinase casein kinase 2 mediates inhibitor-kappaB kinase and aberrant nuclear factor-kappaB activation by serum factor(s) in head and neck squamous carcinoma cells. *Cancer Res*, 2006. 66(13): p. 6722-31.
41. Yu, Y., et al., A novel method for detecting neutralizing antibodies against therapeutic proteins by measuring gene expression. *J Immunol Methods*, 2006. 316(1-2): p. 8-17.
42. Yudt, M.R., et al., Selective and opposing actions of progesterone receptor isoforms in human endometrial stromal cells. *Mol Cell Endocrinol*, 2006. 247(1-2): p. 116-26.
43. Zheng, Z., Y. Luo, and G.K. McMaster, Sensitive and quantitative measurement of gene expression directly from a small amount of whole blood. *Clin Chem*, 2006. 52(7): p. 1294-302.
44. Zimmermann, T.S., et al., RNAi-mediated gene silencing in non-human primates. *Nature*, 2006. 441(7089): p. 111-4.

## 2005

1. Aleksunes, L.M., et al., Differential expression of mouse hepatic transporter genes in response to acetaminophen and carbon tetrachloride. *Toxicol Sci*, 2005. 83(1): p. 44-52.
2. Alikhani, M., Z. Alikhani, and D.T. Graves, FOXO1 functions as a master switch that regulates gene expression necessary for tumor necrosis factor-induced fibroblast apoptosis. *J Biol Chem*, 2005. 280(13): p. 12096-102.
3. Augustine, L.M., et al., Xenobiotic and endobiotic transporter mRNA expression in the blood-testis barrier. *Drug Metab Dispos*, 2005. 33(1): p. 182-9.
4. Burris, T.P., et al., The hypolipidemic natural product guggulsterone is a promiscuous steroid receptor ligand. *Mol Pharmacol*, 2005. 67(3): p. 948-54.
5. Cheng, X., et al., Tissue distribution and ontogeny of mouse organic anion transporting polypeptides (Oatps). *Drug Metab Dispos*, 2005. 33(7): p. 1062-73.
6. Cheng, X., et al., Regulation of mouse organic anion-transporting polypeptides (Oatps) in liver by prototypical microsomal enzyme inducers that activate distinct transcription factor pathways. *Drug Metab Dispos*, 2005. 33(9): p. 1276-82.
7. Crouch, C.R., et al., 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and 1,2,3,4,7,8-hexachlorodibenzo-p-dioxin (HxCDD) alter body weight by decreasing insulin-like growth factor I (IGF-I) signaling. *Toxicol Sci*, 2005. 85(1): p. 560-71.
8. Dai, G., et al., Retinoid X receptor alpha Regulates the expression of glutathione s-transferase genes and modulates acetaminophen-glutathione conjugation in mouse liver. *Mol Pharmacol*, 2005. 68(6): p. 1590-6.
9. Erickson, R.P., et al., Liver disease with altered bile acid transport in Niemann-Pick C mice on a high-fat, 1% cholesterol diet. *Am J Physiol Gastrointest Liver Physiol*, 2005. 289(2): p. G300-7.
10. Gruber, J., et al., RNAi of FACE1 protease results in growth inhibition of human cells expressing lamin A: implications for Hutchinson-Gilford progeria syndrome. *J Cell Sci*, 2005. 118(Pt 4): p. 689-96.
11. Kolachala, V., et al., TNF-alpha upregulates adenosine 2b (A2b) receptor expression and signaling in intestinal epithelial cells: a basis for A2bR overexpression in colitis. *Cell Mol Life Sci*, 2005. 62(22): p. 2647-57.
12. Krishnan, V., et al., The nongenotropic synthetic ligand 4-estren-3alpha17beta-diol is a high-affinity genotropic androgen receptor agonist. *Mol Pharmacol*, 2005. 67(3): p. 744-8.

13. Krishnaswamy, S., et al., UDP glucuronosyltransferase (UGT) 1A6 pharmacogenetics: I. Identification of polymorphisms in the 5'-regulatory and exon 1 regions, and association with human liver UGT1A6 gene expression and glucuronidation. *J Pharmacol Exp Ther*, 2005. 313(3): p. 1331-9.
14. Li, N. and C.D. Klaassen, Lipopolysaccharide-induced down-regulation of organic anion transporting polypeptide 4 (Oatp4; Slc21a10) is independent of tumor necrosis factor-alpha, Interleukin-1beta, interleukin-6, or inducible nitric oxide synthase. *Toxicol Sci*, 2005. 83(1): p. 197-203.
15. Maher, J.M., et al., Induction of the multidrug resistance-associated protein family of transporters by chemical activators of receptor-mediated pathways in mouse liver. *Drug Metab Dispos*, 2005. 33(7): p. 956-62.
16. Maher, J.M., et al., Tissue distribution and hepatic and renal ontogeny of the multidrug resistance-associated protein (Mrp) family in mice. *Drug Metab Dispos*, 2005. 33(7): p. 947-55.
17. Sawada, H., K. Takami, and S. Asahi, A toxicogenomic approach to drug-induced phospholipidosis: analysis of its induction mechanism and establishment of a novel in vitro screening system. *Toxicol Sci*, 2005. 83(2): p. 282-92.
18. Silva, V.M., et al., Transport deficient (TR-) hyperbilirubinemic rats are resistant to acetaminophen hepatotoxicity. *Biochem Pharmacol*, 2005. 70(12): p. 1832-9.
19. Stayrook, K.R., et al., Regulation of carbohydrate metabolism by the farnesoid X receptor. *Endocrinology*, 2005. 146(3): p. 984-91.
20. Thukral, S.K., et al., Prediction of nephrotoxicant action and identification of candidate toxicity-related biomarkers. *Toxicol Pathol*, 2005. 33(3): p. 343-55.
21. Vermeulen, A., et al., The contributions of dsRNA structure to Dicer specificity and efficiency. *Rna*, 2005. 11(5): p. 674-82.
22. Wen, X. and T. Walle, Preferential induction of CYP1B1 by benzo[a]pyrene in human oral epithelial cells: impact on DNA adduct formation and prevention by polyphenols. *Carcinogenesis*, 2005. 26(10): p. 1774-81.
23. Wen, X., U.K. Walle, and T. Walle, 5,7-Dimethoxyflavone downregulates CYP1A1 expression and benzo[a]pyrene-induced DNA binding in Hep G2 cells. *Carcinogenesis*, 2005. 26(4): p. 803-9.

## 2004

1. Buist, S.C. and C.D. Klaassen, Rat and mouse differences in gender-predominant expression of organic anion transporter (Oat1-3; Slc22a6-8) mRNA levels. *Drug Metab Dispos*, 2004. 32(6): p. 620-5.
2. Cherrington, N.J., et al., Lipopolysaccharide-mediated regulation of hepatic transporter mRNA levels in rats. *Drug Metab Dispos*, 2004. 32(7): p. 734-41.
3. Dieter, M.Z., et al., Expression and regulation of the sterol half-transporter genes ABCG5 and ABCG8 in rats. *Comp Biochem Physiol C Toxicol Pharmacol*, 2004. 139(4): p. 209-18.
4. Gruber, J., Functional gene analysis in cultured vertebrate cells using siRNA mediated gene silencing. 2004.
5. Heijne, W.H., et al., Bromobenzene-induced hepatotoxicity at the transcriptome level. *Toxicol Sci*, 2004. 79(2): p. 411-22.
6. Li, N., et al., Down-regulation of mouse organic anion-transporting polypeptide 4 (Oatp4; Oatp1b2; Slc21a10) mRNA by lipopolysaccharide through the toll-like receptor 4 (TLR4). *Drug Metab Dispos*, 2004. 32(11): p. 1265-71.
7. Li, N. and C.D. Klaassen, Role of liver-enriched transcription factors in the down-regulation of organic anion transporting polypeptide 4 (oatp4; oatp1b2; slc21a10) by lipopolysaccharide. *Mol Pharmacol*, 2004. 66(3): p. 694-701.
8. Lu, H., C. Chen, and C. Klaassen, Tissue distribution of concentrative and equilibrative nucleoside transporters in male and female rats and mice. *Drug Metab Dispos*, 2004. 32(12): p. 1455-61.
9. Reynolds, A., et al., Rational siRNA design for RNA interference. *Nat Biotechnol*, 2004. 22(3): p. 326-30.

10. Sakamoto, S., et al., Increased expression of CYR61, an extracellular matrix signaling protein, in human benign prostatic hyperplasia and its regulation by lysophosphatidic acid. *Endocrinology*, 2004. 145(6): p. 2929-40.
11. Soutschek, J., et al., Therapeutic silencing of an endogenous gene by systemic administration of modified siRNAs. *Nature*, 2004. 432(7014): p. 173-8.
12. Wright, S.H., et al., Functional map of TEA transport activity in isolated rabbit renal proximal tubules. *Am J Physiol Renal Physiol*, 2004. 287(3): p. F442-51.

## 2003

1. Bramlett, K.S., et al., A natural product ligand of the oxysterol receptor, liver X receptor. *J Pharmacol Exp Ther*, 2003. 307(1): p. 291-6.
2. Buist, S.C., N.J. Cherrington, and C.D. Klaassen, Endocrine regulation of rat organic anion transporters. *Drug Metab Dispos*, 2003. 31(5): p. 559-64.
3. Cao, G., et al., Antidiabetic action of a liver x receptor agonist mediated by inhibition of hepatic gluconeogenesis. *J Biol Chem*, 2003. 278(2): p. 1131-6.
4. Chen, C., G.E. Hennig, and J.E. Manautou, Hepatobiliary excretion of acetaminophen glutathione conjugate and its derivatives in transport-deficient (TR-) hyperbilirubinemic rats. *Drug Metab Dispos*, 2003. 31(6): p. 798-804.
5. Chen, C., J.L. Staudinger, and C.D. Klaassen, Nuclear receptor, pregnane X receptor, is required for induction of UDP-glucuronosyltransferases in mouse liver by pregnenolone-16 alpha-carbonitrile. *Drug Metab Dispos*, 2003. 31(7): p. 908-15.
6. Cherrington, N.J., et al., Induction of multidrug resistance protein 3 (mrp3) in vivo is independent of constitutive androstane receptor. *Drug Metab Dispos*, 2003. 31(11): p. 1315-9.
7. Choudhuri, S., et al., Constitutive expression of various xenobiotic and endobiotic transporter mRNAs in the choroid plexus of rats. *Drug Metab Dispos*, 2003. 31(11): p. 1337-45.
8. Khvorova, A., A. Reynolds, and S.D. Jayasena, Functional siRNAs and miRNAs exhibit strand bias. *Cell*, 2003. 115(2): p. 209-16.
9. Leazer, T.M. and C.D. Klaassen, The presence of xenobiotic transporters in rat placenta. *Drug Metab Dispos*, 2003. 31(2): p. 153-67.
10. Sakamoto, S., et al., Development of quantitative detection assays for CYR61 as a new marker for benign prostatic hyperplasia. *J Biomol Screen*, 2003. 8(6): p. 701-11.
11. Shelby, M.K., et al., Tissue mRNA expression of the rat UDP-glucuronosyltransferase gene family. *Drug Metab Dispos*, 2003. 31(3): p. 326-33.
12. Slitt, A.L., et al., Induction of multidrug resistance protein 3 in rat liver is associated with altered vectorial excretion of acetaminophen metabolites. *Drug Metab Dispos*, 2003. 31(9): p. 1176-86.
13. Tsai, S.P., et al., Nucleic acid capture assay, a new method for direct quantitation of nucleic acids. *Nucleic Acids Res*, 2003. 31(6): p. e25.
14. Wu, S.J., et al., Quantitation of CYP1A1 and 1B1 mRNA in polycyclic aromatic hydrocarbon-treated human T-47D and HepG2 cells by a modified bDNA assay using fluorescence detection. *Anal Biochem*, 2003. 312(2): p. 162-6.

## 2002

1. Bader, T., et al., Human adipose tissue under in vitro inhibition of 11beta-hydroxysteroid dehydrogenase type 1: differentiation and metabolism changes. *Horm Metab Res*, 2002. 34(11-12): p. 752-7.
2. Brady, J.M., et al., Tissue distribution and chemical induction of multiple drug resistance genes in rats. *Drug Metab Dispos*, 2002. 30(7): p. 838-44.
3. Buist, S.C., et al., Gender-specific and developmental influences on the expression of rat organic anion transporters. *J Pharmacol Exp Ther*, 2002. 301(1): p. 145-51.

4. Cherrington, N.J., et al., Organ distribution of multidrug resistance proteins 1, 2, and 3 (Mrp1, 2, and 3) mRNA and hepatic induction of Mrp3 by constitutive androstane receptor activators in rats. *J Pharmacol Exp Ther*, 2002. 300(1): p. 97-104.
5. Czerwinski, M., et al., Analysis of CYP mRNA expression by branched DNA technology. *Methods Enzymol*, 2002. 357: p. 170-9.
6. Guo, G.L., S. Choudhuri, and C.D. Klaassen, Induction profile of rat organic anion transporting polypeptide 2 (oatp2) by prototypical drug-metabolizing enzyme inducers that activate gene expression through ligand-activated transcription factor pathways. *J Pharmacol Exp Ther*, 2002. 300(1): p. 206-12.
7. Guo, G.L., D.R. Johnson, and C.D. Klaassen, Postnatal expression and induction by pregnenolone-16alpha-carbonitrile of the organic anion-transporting polypeptide 2 in rat liver. *Drug Metab Dispos*, 2002. 30(3): p. 283-8.
8. Harstad, E.B. and C.D. Klaassen, Analysis of strain difference in sensitivity to cadmium-induced hepatotoxicity in Fischer 344 and Sprague-Dawley rats. *Toxicol Sci*, 2002. 67(2): p. 329-40.
9. Johnson, D.R., G.L. Guo, and C.D. Klaassen, Expression of rat Multidrug Resistance Protein 2 (Mrp2) in male and female rats during normal and pregnenolone-16alpha-carbonitrile (PCN)-induced postnatal ontogeny. *Toxicology*, 2002. 178(3): p. 209-19.
10. Johnson, D.R., S.S. Habeebu, and C.D. Klaassen, Increase in bile flow and biliary excretion of glutathione-derived sulfhydryls in rats by drug-metabolizing enzyme inducers is mediated by multidrug resistance protein 2. *Toxicol Sci*, 2002. 66(1): p. 16-26.
11. Johnson, D.R. and C.D. Klaassen, Regulation of rat multidrug resistance protein 2 by classes of prototypical microsomal enzyme inducers that activate distinct transcription pathways. *Toxicol Sci*, 2002. 67(2): p. 182-9.
12. Leazer, T.M., Y. Liu, and C.D. Klaassen, Cadmium absorption and its relationship to divalent metal transporter-1 in the pregnant rat. *Toxicol Appl Pharmacol*, 2002. 185(1): p. 18-24.
13. Li, N., et al., Tissue expression, ontogeny, and inducibility of rat organic anion transporting polypeptide 4. *J Pharmacol Exp Ther*, 2002. 301(2): p. 551-60.
14. Luckey, S.W., et al., 4-hydroxynonenal decreases interleukin-6 expression and protein production in primary rat Kupffer cells by inhibiting nuclear factor-kappaB activation. *J Pharmacol Exp Ther*, 2002. 302(1): p. 296-303.
15. Luo, G., et al., CYP3A4 induction by drugs: correlation between a pregnane X receptor reporter gene assay and CYP3A4 expression in human hepatocytes. *Drug Metab Dispos*, 2002. 30(7): p. 795-804.
16. Park, J.D., N.J. Cherrington, and C.D. Klaassen, Intestinal absorption of cadmium is associated with divalent metal transporter 1 in rats. *Toxicol Sci*, 2002. 68(2): p. 288-94.
17. Park, J.D., S.S. Habeebu, and C.D. Klaassen, Testicular toxicity of di-(2-ethylhexyl)phthalate in young Sprague-Dawley rats. *Toxicology*, 2002. 171(2-3): p. 105-15.
18. Slitt, A.L., et al., Tissue distribution and renal developmental changes in rat organic cation transporter mRNA levels. *Drug Metab Dispos*, 2002. 30(2): p. 212-9.
19. Vansell, N.R. and C.D. Klaassen, Increase in rat liver UDP-glucuronosyltransferase mRNA by microsomal enzyme inducers that enhance thyroid hormone glucuronidation. *Drug Metab Dispos*, 2002. 30(3): p. 240-6.
20. Wagaman, P.C., M.A. Leong, and K.A. Simmen, Development of a novel influenza A antiviral assay. *J Virol Methods*, 2002. 105(1): p. 105-14.
21. Xiang, H., et al., Enhanced tumor killing by Apo2L/TRAIL and CPT-11 co-treatment is associated with p21 cleavage and differential regulation of Apo2L/TRAIL ligand and its receptors. *Oncogene*, 2002. 21(22): p. 3611-9.
22. Zhang, Y., et al., Liver X receptor and retinoic X receptor mediated ABCA1 regulation and cholesterol efflux in macrophage cells-messenger RNA measured by branched DNA technology. *Mol Genet Metab*, 2002. 77(1-2): p. 150-8.

## 2001

1. Allan, G.F., et al., Induction of the progesterone receptor gene in estrogen target cells monitored by branched DNA signal amplification. *Steroids*, 2001. 66(9): p. 663-71.
2. Player, A.N., et al., Single-copy gene detection using branched DNA (bDNA) in situ hybridization. *J Histochem Cytochem*, 2001. 49(5): p. 603-12.
3. Rausch-Derra, L.C., et al., Differential effects of microsomal enzyme-inducing chemicals on the hepatic expression of rat organic anion transporters, OATP1 and OATP2. *Hepatology*, 2001. 33(6): p. 1469-78.
4. Sen, A., et al., Adipogenic potential of human adipose derived stromal cells from multiple donors is heterogeneous. *J Cell Biochem*, 2001. 81(2): p. 312-9.
5. Takiguchi, M., et al., Cyproterone acetate induces a cellular tolerance to cadmium in rat liver epithelial cells involving reduced cadmium accumulation. *Toxicology*, 2001. 165(1): p. 13-25.

## 2000

1. Bramlett, K.S., S. Yao, and T.P. Burris, Correlation of farnesoid X receptor coactivator recruitment and cholesterol 7alpha-hydroxylase gene repression by bile acids. *Mol Genet Metab*, 2000. 71(4): p. 609-15.
2. Hartley, D.P. and C.D. Klaassen, Detection of chemical-induced differential expression of rat hepatic cytochrome P450 mRNA transcripts using branched DNA signal amplification technology. *Drug Metab Dispos*, 2000. 28(5): p. 608-16.
3. Warrior, U., et al., Application of QuantiGene nucleic acid quantification technology for high throughput screening. *J Biomol Screen*, 2000. 5(5): p. 343-52.
4. Wentworth, J.M., T.P. Burris, and V.K. Chatterjee, HIV protease inhibitors block human preadipocyte differentiation, but not via the PPARgamma/RXR heterodimer. *J Endocrinol*, 2000. 164(2): p. R7-R10.
5. Zhou, L., et al., A branched DNA signal amplification assay to quantitate messenger RNA of human uncoupling proteins 1, 2, and 3. *Anal Biochem*, 2000. 282(1): p. 46-53.

## 1999

1. Burris, T.P., et al., A novel method for analysis of nuclear receptor function at natural promoters: peroxisome proliferator-activated receptor gamma agonist actions on aP2 gene expression detected using branched DNA messenger RNA quantitation. *Mol Endocrinol*, 1999. 13(3): p. 410-7.
2. Todd, M.D., et al., Toxicity Screening of a Combinatorial Library: Correlation of Cytotoxicity and Gene Induction to Compound Structure. *J Biomol Screen*, 1999. 4(5): p. 259-268.